

Prevention and Schizophrenia—The Role of Dietary Factors

John McGrath^{*,1,2}, Alan Brown³, and David St Clair⁴

¹Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, Australia; ²Queensland Brain Institute and Department of Psychiatry, University of Queensland, St Lucia QLD 4072, Australia; ³College of Physicians and Surgeons of Columbia University, New York State Psychiatric Institute, New York, NY; ⁴Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen, UK

*To whom correspondence should be addressed; tel: +61-7-3271-8694, fax: +61-7-3271-8698, e-mail: john_mcgrath@qcmhr.uq.edu

Adequate prenatal nutrition is essential for optimal brain development. There is a growing body of evidence from epidemiology linking exposure to nutritional deprivation and increased risk of schizophrenia. Based on studies from the Netherlands and China, those exposed to macronutrient deficiencies during famine have an increased risk of schizophrenia. With respect to micronutrients, we focus on 3 candidates where there is biological plausibility for a role in this disorder and at least 1 study of an association with schizophrenia. These nutrients include vitamin D, folic acid, and iron. While the current evidence is incomplete, we discuss the potential implications of these findings for the prevention of schizophrenia. We argue that schizophrenia can draw inspiration from public health interventions related to prenatal nutrition and other outcomes and speculate on relevant factors that bear on the nature, risks, impact, and logistics of various nutritional strategies that may be employed to prevent this disorder.

Key words: nutrition/epidemiology/prevention/schizophrenia

Introduction

Nutritional deprivation (ND) represents one of the greatest public health challenges throughout the world. The prevalence of maternal undernutrition varies from 10% to 19% in most countries, although the most deprived countries, such as those in South Asia and Africa, experience a prevalence of approximately 40%.¹ Subjects who are nutritionally deprived may suffer from both macro- and micronutrient deficiencies. Although macronutrient deficiency is not common in high-income countries, where food is more plentiful and obesity is more of a problem, deficiencies of micronutrients have a surprisingly high prevalence.

Pregnancy is a time of increased nutrient demand in order to satisfy normal fetal growth and development,

resulting in an even higher prevalence of ND in this population, and, as reviewed below, ND during pregnancy is a major cause of disease and disability in the offspring. Hence, there is a pressing need for preventive nutritional interventions during pregnancy.

ND during gestation and infancy represents one of the most common causes of neurodevelopmental disorders, and, as reviewed elsewhere, schizophrenia is likely rooted in deviations in neurodevelopment.^{2,3} Hence, the role of prenatal ND has been investigated as a potential etiologic factor in schizophrenia. A major goal of this work is to assess the extent to which schizophrenia might be preventable by nutritional interventions during gestation.

In this article, we provide brief overviews of the evidence supporting associations between macro- and micronutrient deficiency in schizophrenia. We focus on 3 micronutrients in which there is biological plausibility for a role in this disorder and at least 1 study of an association with schizophrenia. These nutrients include vitamin D, folic acid, and iron. We then discuss the potential implications of these findings for the prevention of schizophrenia. Where possible, we shall report the population attributable risk (PAR) (see Brown and McGrath⁴ for a detailed discussion about the limitations of this metric). Drawing on lessons learned from nutritional interventions in other disorders, and the extant biological literature, we shall speculate on relevant factors that bear on the nature, risks, impact, and logistics of various nutritional strategies that may be employed to prevent this disorder.

Studies of Prenatal Exposure to Famine And General Indicators of Malnutrition

In general, if prenatal malnutrition was associated with an increased risk of schizophrenia, then one would predict that the incidence of schizophrenia should be higher

in countries with endemic malnutrition. However, there is a lack of high-quality incidence data from low-income countries.⁵ Furthermore, a wide range of epidemiological factors complicate the interpretation of simple ecological analyses of this research question (eg, differential infant and childhood mortality, the influence of comorbid neurodevelopmental disorders, between-site differences in putative protective factors related to rural settings, and cohesive cultural influences). However, in recent years more precise epidemiological studies have been able to explore the association between prenatal exposure to famine and risk of schizophrenia.

The first direct evidence that prenatal ND is related to schizophrenia derived from studies of the Dutch Hunger Winter of 1944–1945, in which a Nazi blockade of the Western Netherlands resulted in an acute period of starvation for most of the inhabitants of that region.⁶ Mortality increased 2-fold, and fertility was halved. This was a severe tragedy, though from it came a series of studies on the role of prenatal starvation in health outcomes, including psychiatric disorders. Three key elements made this research possible: precise documentation of caloric food rations, a well-demarcated period of severe famine, and a comprehensive nationwide psychiatric registry. Equipped with these design features, Susser et al^{7,8} demonstrated that exposure to severe famine was associated with a 2-fold increase in risk of schizophrenia, coinciding with a peak in risk of Central Nervous System anomalies, including neural tube defects (NTDs). Over a decade later, St Clair and colleagues replicated this finding in 2 regions of China, which experienced a famine during 1959–1961, a result of Mao Zedong's Great Leap Forward.^{9,10} Although the Chinese studies did not have precise data on the timing of the exposures, the sample sizes were 10–15 times those in the Dutch study, and the effect sizes found in both Chinese studies were similar to the original Dutch results. Overall, the 3 studies provide strong evidence that prenatal ND plays a role in schizophrenia.

Clearly, severe famine is associated with catastrophic health outcomes for pregnant women and their offspring. The 3 famine studies suggest that offspring who survive severe prenatal restrictions in proteins, calories, and micronutrients have altered brain development that increases their risk of later schizophrenia.

Prenatal Micronutrients And Schizophrenia

Compared with ecological studies related to macronutrition, investigations that document levels of specific micronutrients are likely to offer more promise with respect to the development of preventive strategies for schizophrenia because of the solid body of public health research related to prenatal nutrition and infant outcomes. Although this work is only in its infancy, there are already several intriguing suggestions that prenatal deficiencies of micronutrients are implicated in schizo-

phrenia among offspring. Below, we briefly review these findings.

Vitamin D

The vitamin D₃ system is unique in the sense that its production depends primarily on the action of sunlight on the skin.¹¹ Ultraviolet B radiation acts on a cholesterol metabolite in the epidermis to produce previtamin D. Subsequent hydroxylations in the liver and kidney create the active moiety (which is a seco-steroid hormone). Vitamin D production is associated with the duration of the photoperiod, which in turn is influenced by latitude and season.^{11,12} A minor source of vitamin D is via the diet because certain foods contain vitamin D (eg, fatty fish). Vitamin D can also be obtained via supplements or the consumption of fortified foods (eg, some dairy products). From a global perspective, hypovitaminosis D is one of the most prevalent micronutrient deficiencies.¹³

In order to explore the biological plausibility of low prenatal vitamin D as a risk factor for schizophrenia, an animal model has been developed in which female rats are depleted of vitamin D prior to mating and throughout pregnancy but return to normal diet after the litter is born. This model is called the Developmental Vitamin D (DVD) deficiency model. Neurobiological findings from these animals include larger lateral ventricles, increased cellular proliferation, reduced apoptosis, altered neurogenesis,^{14–16} persistent ventricular enlargement,¹⁷ hyperlocomotion in novel environments,^{18,19} enhanced locomotion in response to psychomimetic agents, such as the *N*-methyl-D-aspartate antagonist MK-801^{19,20} and amphetamine,²¹ increased sensitivity to haloperidol,¹⁹ and altered attentional processing (impaired latent inhibition,²² altered learning,²³ and altered synaptic plasticity²⁴).

Epidemiological Evidence of Vitamin D Deficiency And Schizophrenia

There is growing recognition that vitamin D deficiency is prevalent in many nations.^{25,26} Pregnant women are also vulnerable to hypovitaminosis D, due to alterations in outdoor behavior and increased needs of the developing fetus.²⁷ Ecological studies support a potential role for vitamin D in schizophrenia.²⁸ These data include studies that have explored the association between schizophrenia and winter/spring birth²⁹ and also the apparent increased incidence and prevalence of schizophrenia at higher latitudes.^{30–32} An additional supportive finding is the observation that dark-skinned migrant groups, who have a higher prevalence of hypovitaminosis D, have an increased risk of psychosis.^{5,33,34} Vitamin D deficiency may account for a proportion of the increased risk associated with migrant status.³⁵

While the clues from ecological studies make vitamin D an attractive candidate, to date the evidence from

analytical epidemiology is modest. In the first attempt to test this hypothesis, a small pilot study based on 26 cases and 51 controls examined the storage form of vitamin D (25-hydroxyvitamin D) in third trimester maternal sera³⁶; however, the results of this pilot study were inconclusive. Based on the Northern Finnish birth cohort study, a link between the absence of vitamin D supplementation during the first year of life and a significantly increased risk of schizophrenia in men was found.³⁷ Recently, a study based on Danish case-control study examined vitamin D status in neonatal dried blood spots.³⁸ As expected, the study confirmed that 25-hydroxyvitamin D concentrations showed significant seasonal variation and were significantly lower in the offspring of migrants compared with native-born parents. Overall, the risk of schizophrenia was significantly associated with the level of vitamin D, with lower levels of vitamin D associated with an increased risk of schizophrenia compared with the reference category (fourth quintile). However, the relationship was nonlinear, with the highest quintile of vitamin D also associated with a slightly increased risk of schizophrenia. The PAR for vitamin D in this study was 43.6%. The findings require replication, and the mechanisms underlying the nonlinear association remain to be clarified.

Folic Acid And Homocysteine

As discussed above, a sharp increase in prevalence of both schizophrenia and NTDs was related to periconceptional and early gestational exposure to the Dutch Hunger Winter.³⁹ A wealth of data suggest that folic acid supplementation, either through vitamins or fortified grains,⁴⁰ leads to a marked decline in incidence of NTDs, suggesting that a deficit of this nutrient might be responsible for both schizophrenia and NTDs.

The relationship between maternal levels of homocysteine, an amino acid central to the folate metabolic cycle, and risk of schizophrenia in offspring has been investigated. Although folate levels degrade in stored sera, homocysteine represents an excellent proxy given that levels of this amino acid reliably increase when folate levels decline. Moreover, homocysteine acts as a partial antagonist on the *N*-methyl-D-aspartate receptor (NMDAR) at physiologic glycine concentrations,⁴¹ consistent with evidence for NMDAR hypofunction in schizophrenia, and hyperhomocysteinemia disrupts placental function, through vasculopathic and apoptotic effects,^{42,43} which may compromise oxygen delivery to the fetus. Homocysteine also alters levels of the methyl donor S-adenosylmethionine (S-AdoMet), which regulates the expression of many genes involved in neurodevelopment.⁴⁴

Hence, the relationship between maternal homocysteine and offspring risk of schizophrenia was investigated in the Child Health and Development Study (CHDS) birth cohort in northern California, which

had available archived sera drawn during pregnancy in nearly all members of this cohort. Elevated third trimester maternal homocysteine levels were associated with a greater than 2-fold increase in the risk of schizophrenia.⁴⁵

Iron

Iron deficiency anemia (IDA) is one of the most common nutrient deficiencies in pregnant women.⁴⁶ IDA is defined as anemia accompanied by depleted stores of iron and signs of compromised iron supply to the tissues. The main etiologies are inadequate dietary intake, an increased requirement for iron, parasitic infections, and loss of blood from other causes.⁴⁷ With respect to the global burden of disease, those with tropical diseases are at increased risk of iron deficiency.⁴⁸ The global prevalence of IDA during pregnancy is 42%.⁴⁹ Pregnancy is known to increase the demands for iron due to expansion of the maternal and fetal blood supply and growth and development of the fetus.

Iron is essential for a plethora of metabolic processes that are essential for optimal brain function. These include its role as a coenzyme of dopamine synthesis and its relationship to dopamine receptor density⁵⁰; dopaminergic abnormalities are a hallmark feature of schizophrenia.^{51–53} Iron is also an essential cofactor for synthesis of lipids and cholesterol; this explains disruptions of myelination following prenatal iron deficiency,⁵⁴ and these deficits did not improve despite restoration of iron at weaning.^{55,56} Intriguingly, myelin and oligodendrocyte abnormalities have been demonstrated in patients with schizophrenia.^{57–59} Perinatal iron deficiency also alters the timing and expression of genes that are necessary for development of the hippocampus.⁶⁰ Animal experiments have confirmed that pre- and perinatal iron deficiencies are associated with a range of neurobiological outcomes of interest to schizophrenia (eg, working memory, hippocampal size, gene expression, and synaptic plasticity).^{61–63} IDA is a risk factor for adverse pregnancy outcomes, including fetal hypoxia, intrauterine growth retardation, and prematurity.^{64–68} Moreover, iron deficiency during pregnancy, as well as in infancy and childhood, has long been associated with motor, cognitive, and behavioral abnormalities^{69–71} similar to those observed in children who later developed schizophrenia.^{72–74}

These suggestions led to the first study specifically aimed at evaluating the role of maternal iron deficiency in schizophrenia. In the CHDS cohort, a mean maternal hemoglobin concentration of 10 g/dl, indicative of maternal anemia, was associated with a nearly 4-fold increased rate of developing schizophrenia, adjusting for maternal education and ethnicity.⁷⁵ A dose-response relationship was observed, with a graded increase in risk of schizophrenia correlating with increasing hemoglobin levels.

This association has been recently replicated in a large Danish population-based cohort.⁷⁶

Primary Preventive Approaches

While there remains a paucity of data supporting a role for vitamin D, folic acid, and iron in the etiology of schizophrenia, it is worth noting that the findings reported above on these micronutrients are all based on prospectively acquired data from maternal or neonatal biomarkers and comprehensive follow-up of the offspring for schizophrenia. Nonetheless, of these 3 micronutrients, an attempt to replicate a previous association has been conducted only for iron. Hence, the following discussion on primary prevention of schizophrenia through micronutrient supplementation should be viewed with appropriate caution.

Documenting an association between a prenatal nutrient of potential causal relevance to a disorder in the offspring, though of high importance, is only the first step toward the development of preventive approaches. In our view, implementing a prophylactic nutritional intervention for schizophrenia should be predicated on 4 additional factors. First, what are the types of nutritional interventions that can be implemented, and what is the evidence that such interventions are effective in preventing adverse outcomes during infancy and childhood? Second, what is the “optimum” quantity of intake of such nutrients in pregnancy, when in pregnancy should they be taken, and are there any risks of such supplementation? Third, if the associations between micronutrient deficiency and schizophrenia are confirmed, what is the potential impact of correcting the nutritional intervention on reducing risk of schizophrenia? Fourth, what are the logistics and practicality of such interventions and relative costs for preventing schizophrenia for each of these 3 nutrients? To address these questions, we shall consider lessons learned from nutritional interventions aimed at the primary prevention of other reproductive outcomes.

Vitamin D

With respect to the potential links between vitamin D and schizophrenia, more work needs to be done before any recommendation can be made. Worryingly, there is evidence that the relationship between prenatal vitamin D and neonatal outcomes⁷⁷ and schizophrenia³⁸ is nonlinear. Until the nature of this relationship is resolved, simple public health recommendations are difficult to formulate. In some respects, if the evidence mounts that neonatal and childhood outcomes like bone density can be improved with prenatal vitamin D supplements, this type of evidence is more likely to influence public health bodies to implement more assertive maternal and neonatal supplementation programs. There may

be opportunities for “natural experiments” to explore these issues, much as has been done with interventions related to perinatal folate supplementation and spina bifida or neonatal sleeping position and Sudden Infant Death syndrome.

While the focus of research related to vitamin D and schizophrenia has focused on prenatal exposures, it is feasible that hypovitaminosis D during childhood and adolescence could also impact on mental health.⁷⁸ Second-generation dark-skinned migrants to cold climates are exposed to low vitamin D across the life span, thus cumulative prenatal and postnatal exposures may be additive with respect to subsequent risk of psychosis. With respect to first-generation migrants, we lack research that has examined age-at-first-migration and subsequent risk of psychosis. Brain growth is extremely rapid in the first few years of life, thus it is feasible that the absence of vitamin D could disrupt cell cycle kinetics and a range of neurotropic factors.⁷⁹ Puberty may also be a window of vulnerability because sex hormones and vitamin D share a broad range of downstream coactivating agents. With respect to prevention, it is possible that supplementation during childhood and adolescence could avert conversion to schizophrenia. While speculative, there is a cogent argument to fast track any type of safe intervention that could possibly result in the primary prevention of schizophrenia. With respect to the potential primary prevention of schizophrenia, vitamin D seems to be an attractive candidate from a public health perspective. It is a cheap, safe, and relatively simple intervention that could potentially avert a wide range of disorders other than schizophrenia.^{80,81}

Iron

Several interventions have been employed in previous studies of iron during pregnancy and reproductive outcomes, including iron supplementation via tablets, iron fortification of food, education on health and nutrition, control of parasitic infections, and improved sanitation.⁸² Among these, iron supplementation has been most commonly used. These studies have demonstrated that this intervention leads to a substantial reduction of anemia in late pregnancy, at delivery, and 6 weeks postpartum.^{83,84}

Iron supplementation appears to lead to favorable perinatal outcomes. Pregnant women with evidence of anemia who were supplemented with daily iron, compared with no/placebo intervention, had an approximately 20% diminished likelihood of having offspring with low birth weight, and birth length was increased by 0.38 cm.⁸⁵ However, no differences were found in these outcomes when women were not selected for anemia status and for a number of additional outcomes, including prematurity, placental abruption, preeclampsia, or puerperal infection. Infants whose mothers received

a combination of daily iron and folic acid during pregnancy were 57.7 g heavier than infants from pregnancies without this supplementation^{86,87} and were less likely to give birth to offspring of small gestational age.⁸⁶

Data on maternal iron supplementation during pregnancy and neurocognitive and behavioral development in offspring are more mixed. In a systematic review and meta-analysis of 17 randomized controlled trials (RCTs), a modest improvement in IQ in children aged >7 years was observed; however, no effect was observed on the Bayley Mental Development Index and motor development score.⁸⁸ In a later study of offspring at age 4, maternal iron supplementation was not associated with improved IQ or behavior,⁸⁹ and another study of offspring at age 6–8 indicated no association with mean behavior and temperament scores, although surprisingly peer problems were reported to be greater in the offspring of mothers who received iron during pregnancy than placebo.⁹⁰

While the findings do not show a clear effect of iron supplementation on cognitive and behavioral outcomes, the lack of consistent results may be accounted for by limitations that could be instructive for the evaluation of preventive approaches for schizophrenia. First, all the interventions were applied to general populations rather than targeted for women with IDA. It is possible that if the interventions were targeted to women suffering from anemia, positive effects on child cognitive and behavioral outcomes may have been observed, as was found for birth weight and birth length among mothers who were supplemented with iron in the studies mentioned above.⁸⁵ This could be of particular relevance to low-income settings. Hence, although more trials are needed on both anemic and nonanemic populations, these data raise the question of whether efforts should be made to target pregnancies that are vulnerable to iron deficiency. Second, the studies varied by the quantities of iron administered. This raises the question as to the “optimum” quantity of iron to be administered. Current recommendations vary, from 27 mg/day (by the Institute of Medicine) to as much as 60–120 mg/day,⁹¹ depending on how many months of treatment can be achieved during pregnancy, the presence of anemia in the mother, and the prevalence of iron deficiency or anemia in the population. Yet, these recommendations are not based on the dosage of iron necessary to prevent neuropsychiatric outcomes but rather on that needed to prevent anemia in the mother and newborn. The upper limit of iron is based largely on the potential for detrimental side effects, including gastrointestinal consequences such as diarrhea, constipation, and abdominal discomfort,⁹² particularly at the higher doses. However, a recent study of iron prophylaxis in pregnancy did not demonstrate any increase in side effects at doses up to 80 mg/day.⁹³ Third, differences in the period of administration during pregnancy may have accounted for inconsistent results. For schizophre-

nia, associations with iron deficiency were present for both the second and the third trimesters,⁷⁵ though clearly more studies will be necessary to confirm that there is a critical “window” of supplementation necessary to prevent this outcome.

Given the lack of epidemiological studies, and the need for more data on the effects of iron supplementation on neuropsychiatric outcomes, we can only speculate on the potential impact of this intervention. Based on the first study of maternal iron and schizophrenia cited,⁷⁵ the PAR is 11.6%. These findings suggest that iron supplementation could play an appreciable role in reducing risk of schizophrenia.

Is iron supplementation a practical intervention? A recent randomized controlled trial suggests that approximately 17% of pregnant women given oral iron during pregnancy did not continue iron due mostly to side effects and that 12.5% were noncompliant.⁹² In another study of iron supplementation, in rural China, however, compliance was high, with supplements being consumed on greater than 90% of the days on which they were available. Nonetheless, there is a lack of good studies on how well adherence to an iron supplementation regimen would fare in an obstetric practice, and there would clearly be differences depending upon the context, ie, industrialized vs developing country, urban vs rural population, high vs low income, and type of treatment setting.

With regard to cost, iron is remarkably affordable, particularly in comparison with prescribed medications. In a recent study, the cost of 1 iron sulfate tablet (250 mg) was \$0.5 (United States), for an average cost for an entire pregnancy of \$165.⁹² Although there are insufficient data to estimate the potential preventive effect of a prenatal iron supplementation program on schizophrenia risk, this calculation suggests that this intervention is equally to or more affordable than many current diagnostic and therapeutic interventions in obstetric practice.

However, for the billions of individuals who exist on less than \$1.25 dollars per day, iron supplementation during pregnancy is unaffordable in the absence of funded public health care programs. Most of the anemia in this group is a consequence of common tropical diseases (eg, hookworm affects over 500 million people). These disorders are amenable to cost-effective coordinated public health packages.^{94,95}

Folate

Folic acid supplementation in the prevention of NTDs is one of the major public health success stories of the 20th century. In a recent review by the US Preventive Services Task Force of several large-scale studies, ORs for reductions of NTDs following periconceptional folic acid supplementation ranged from 0.11 to 0.65, indicating a reduction in risk of NTDs between 35% and 89%.⁴⁰ These studies included cohort and case-control studies

and RCTs. Periconceptional use of multivitamins containing folic acid have also been demonstrated to decrease the risk of other birth defects, including congenital cardiac anomalies and orofacial clefts.^{96,97} These studies included folate prescribed either as part of a multivitamin or as folic acid tablets. Hence, periconceptional folic acid administration has clearly been shown to be of benefit in preventing developmental outcomes, though no known studies have examined folic acid supplementation in relation to neuropsychiatric outcomes other than NTDs.

In 1998, the US Food and Drug Administration introduced a universal prevention program consisting of mandated folic acid fortification (FAF) of enriched cereal-grain products. This represented a sharp departure from the previous policy of targeting pregnant populations with folate supplementation. In an analysis by the Centers for Disease Control and Prevention (CDC) of births from 23 states and Puerto Rico before and after the introduction of FAF, the prevalence of NTDs was reduced by 26%–27%.⁹⁸ In Canada, which also instituted FAF in 1998, a 46% reduction of NTDs was reported.⁹⁹ Mandated fortification of folic acid in the United States has been related to modest but statistically significant decreases in several other birth defects.¹⁰⁰ Most, but not all, developed nations have mandatory folate supplementation in various food products.¹⁰¹

The “optimum” dose of folic acid has yet to be defined. In previous studies of folic acid given as supplements, quantities varied from 0.2 to 0.8 mg/day. In studies of FAF of grains, the recommended US Food and Drug Administration level was 140 µg/100 g,¹⁰² while in Canada fortification was 150 µg/100 g. These amounts were well below those recommended by several established organizations including the CDC, the American Medical Association, and the American College of Obstetricians and Gynecologists, which argued for a substantially greater level of fortification (350 µg/100 g) based on models predicting that the mandated dose would only reduce the rate of NTDs by 20%.^{102–104} As noted above, the empirical data exceeded these estimates. However, in a double-blind “dose finding” trial of supplemental folate in the prevention of NTDs, 200 µg/day of folate provided equal effectiveness to 400 µg/day, but 100 µg/day was insufficient.

More recent studies have investigated the role of plasma folate and of folic acid supplements and childhood neurocognition and behavior. Maternal plasma folate was positively associated with children’s cognitive test scores in domains that included learning, memory, visuospatial ability, and concentration.¹⁰⁵ In a second study, however, no associations between maternal folate and childhood neurocognitive measures were demonstrated.¹⁰⁶ Folic acid supplementation during pregnancy, obtained prospectively in a population-based birth cohort, was associated with greater verbal, motor, executive

function, and social competence.¹⁰⁷ Maternal dietary folate was associated in another study with improved mental development in children, but the effect was isolated to children of mothers who were homozygous for a mutation in the gene for methylenetetrahydrofolate reductase (*MTHFR*),¹⁰⁸ which plays an important role in folate metabolism.

Until recently, folate supplementation was considered to have primarily beneficial effects. Indeed, folate has been shown to maintain genome stability through regulation of DNA biosynthesis, repair, and methylation, and folate deficiency is known to induce and accelerate carcinogenesis by disrupting these processes.¹⁰⁹ This includes its role in remethylating homocysteine to methionine, which is metabolized to SAMe, the principal methyl donor, which regulates gene transcription and protein expression by methylating cytosine. Folate is also necessary for the synthesis of both purines and pyrimidine nucleoside thymidine; folate deficiency, by disrupting the balance between DNA precursors, results in abnormal DNA repair and leads to chromosomal damage by misincorporation of uracil in place of thymidine. This process is also associated with increased risk of malignant transformation. Folate supplementation has been associated in retrospective, case-control, and prospective studies with diminished risk of colorectal cancer.¹¹⁰

However, evidence is emerging that long-term intervention with folic acid may promote growth of initiated cancer cells.^{111,112} This may increase the acceleration of carcinogenesis if folic acid is administered following the emergence of precancerous lesions and subclinical cancers, possibly by providing DNA precursors for growth of cancer cells.¹¹³ In contrast to the previous studies that demonstrated a protective effect on cancer, some observational and controlled folate intervention trials suggest increases in risk of breast, lung, and prostate cancers.¹¹⁴ Hence, while folate may protect against cancer initiation, there is the potential for this vitamin to facilitate progression and growth of preneoplastic lesions and subclinical cancers. It is also possible, though unproven, that increased methylation of DNA cytosine residues could silence promoter activity, thereby inactivating tumor-suppressor genes or other genes that inhibit carcinogenesis.¹¹⁵

With regard to periconceptional folate supplementation, there is recent evidence that this vitamin leads to increased methylation of the maternally imprinted insulin-like growth factor 2 (*IGF2*) gene, possibly due to a relative intrauterine silencing of *IGF2*.¹¹⁶ An increase in folate intake also has the potential to delay the diagnosis of vitamin B12 deficiency, by correcting the macrocytosis,¹¹⁷ and to decrease natural killer cell activity from circulating unmetabolized folic acid,¹¹⁸ though to date no untoward effects of the latter process have been demonstrated. Recently, evidence has emerged linking prenatal

folate supplementation and an increased risk of childhood asthma in the offspring.^{119,120}

It should be kept in mind that in most trials of folate supplementation the vitamin is administered to women planning a pregnancy, in order to ensure that periconceptional levels of folate are adequate, but in some studies folate is continued only until the third month of pregnancy. Yet, as noted previously, in the only study to examine the relationship between a biomarker related to folate status—elevated homocysteine—the association was observed only for the third trimester.¹²¹ Universal fortification of folic acid may be helping to address this issue.

With regard to the potential impact of folic acid interventions during pregnancy, we calculated the PAR for elevated homocysteine and schizophrenia. These findings suggest that if folic acid interventions reduced all homocysteine levels to the lowest 2 tertiles of the distribution, the risk of schizophrenia would decline by 29.6% (PAR). Hence, if the association between elevated maternal homocysteine and schizophrenia risk can be replicated, and found to be of similar magnitude, this suggests that folic acid supplementation and/or fortification strategies may have a considerable impact on reduction of schizophrenia risk in the population.

Unlike iron supplementation, the use of universal FAF does not present the same logistical issues with regard to administration of, and compliance with, this nutrient during pregnancy. However, it is unclear whether the FAF initiative implemented over 10 years ago will provide an adequate amount of folic acid to attain an appreciable decrease in schizophrenia risk over the coming decades. In light of this, 1 reasonable option, at least until clearly harmful effects of universal FAF can be demonstrated, would be to continue the current recommendations for folic acid supplementation during pregnancy, even though these women are already receiving universal FAF through the diet. Consideration might also be given to targeted interventions in pregnancies of women with lifestyle, demographic, environmental, and genetic factors that predispose to low folate intake or low blood folate concentrations. These include, eg, smoking, substance use or abuse, obesity, malabsorption diseases, use of antiepileptic medications, folic acid antagonists, MTHFR genotype (a known polymorphism of which results in diminished folate levels), personal or family history of NTD, and maternal ethnicity (northern Chinese, native American, and Sikh).¹¹⁷

The annual cost of folic acid supplementation or fortification is as low as \$12 per pregnancy, over 10 times less than that of iron. Previous cost-benefit analyses of folic acid revealed a net savings of \$92 000 for low-level fortification per NTD prevented and \$132 000 for folate supplementation per NTD prevented. If lowering homocysteine levels has an appreciable effect of reducing risk of schizophrenia, the cost:benefit ratio for folic acid

interventions may be considerably higher, given that schizophrenia is a far more common outcome.

Conclusion

Evidence from ecological studies has accumulated demonstrating an association with prenatal ND and increased risk of schizophrenia. Other unknown factors must, however, operate alongside micronutrient deficiency; otherwise, we would expect to increased rates of schizophrenia in developing countries compared with advanced economies. While there is a lack of quality epidemiological data on these issues,⁵ this does not seem to be the case. It is our view that before preventative approaches can be considered, well-conducted observational epidemiological studies and robust biological evidence supporting the plausibility of specific nutritional risk factors are essential. Nevertheless, although the data are sparse, and while we lack data from randomized controlled trials, initial birth cohort studies with prospectively acquired data suggest that deficiencies of 3 micronutrients that are biologically relevant to pathogenic processes in schizophrenia are associated with an elevated risk of this disorder.

In keeping with findings from nutritional epidemiology in general, some of the findings suggest a nonlinear relationship between the candidate nutritional agent and risk of schizophrenia. As has been found with birth weight and infant mortality, both very high and very low levels can be associated with risk. The gold standard evidence will require randomized controlled trials of prenatal nutritional supplementation and prospective evaluation of mental health outcomes in the offspring. Because the incidence of schizophrenia is relatively low and because the peak age of onset is in the second and third decade of life, these trials will be a challenge. In the decades to come, the research community should be able to explore the associations between prenatal nutrition and offspring mental health using large prospectively collected cohort of mothers and babies (eg, the Norwegian Mother and Child Cohort,¹²² the US National Children's Study¹²³). Caution is needed when pooling data from different nations because populations from these sites may vary on a wide range of factors (eg, patterns of general nutrition, socioeconomic status, and ethnicity). Thus, the findings from 1 study may not generalize to all populations.

The falling costs of genotyping will also facilitate studies that explore the genetic architecture underpinning candidate micronutritional pathways (eg, polymorphisms that are associated with differential sensitivity to iron, folate, or vitamin D concentrations).^{124–126} These clues can contribute to future gene by environment studies, where prenatal nutrition can be included in models that examine maternal and/or offspring genotype.¹²⁷

While one may argue that it is premature to think about preventative nutrient strategies given the relative

paucity of data, in our opinion, such consideration is necessary for 3 reasons. First, an awareness of the merits of nutritional supplementation and food fortification campaigns in the prevention of other disorders may inspire the research community to recognize the importance of conducting prospective studies of prenatal nutrients in schizophrenia in existing cohorts. For example, there is good evidence that prenatal vitamin D supplements can improve a range of bone-related neonatal outcomes.⁸¹ If randomized controlled trials of various prenatal nutritional interventions are commenced, we recommend that neonatal cognitive outcomes be included. Ideally, if feasible, these cohorts could be followed up for later mental illness outcomes. Folic acid supplementation and fortification has played a critical role in reducing risk of NTDs (and possibly other childhood neurodevelopmental outcomes), and iron supplementation has diminished the risk of certain reproductive outcomes. Second, viewing ND and schizophrenia through the lens of a preventative approach may provide researchers with a new tool to evaluate the utility of investigating certain types of micronutrient or other NDs. Considerations in this regard include the prevalence of the exposure in the population (which is probably the most significant determinant of the PAR and number needed to prevent), whether subpopulations exist that may especially benefit from the intervention, whether the intervention can be practically applied in a public health program or obstetric practice, possible adverse effects of the exposure, and affordability as a public health intervention. Finally, if the associations between ND and schizophrenia are confirmed, considering ND from the standpoint of preventative approaches may have utility in the development of public health policies that can be implemented in the near future in order to reduce the incidence of schizophrenia decades from now.

Funding

National Institute of Mental Health (K02 MH065422-06 to A.B.); Australian National Health and Medical Research Council (J.M.).

Acknowledgments

The authors wish to acknowledge funding support from National Institute of Mental Health and the Australian National Health and Medical Research Council and Patric Prado for his assistance. Drs Brown, McGrath, and St Clair have no conflicts of interest to disclose.

References

- Black R, Allen L, Bhutta Z, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*. 2008;371:243–260.
- Weinberger DR. Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry*. 1987;44:660–669.
- Lewis SW, Murray RM. Obstetric complications, neurodevelopmental deviance, and risk of schizophrenia. *J Psychiatr Res*. 1987;21:413–421.
- Brown AS, Patterson PH. Maternal Infection and Schizophrenia: Implications for Prevention. *Schizophrenia Bulletin*. In press.
- McGrath J, Saha S, Welham J, El Saadi O, MacCauley C, Chant D. A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology. *BMC Med*. 2004;2(1):13.
- Stein Z, Susser M, Saenger G, Marolla F. *Famine and Human Development. The Dutch Hunger Winter of 1944–1945*. London: New York Oxford University Press; 1975.
- Susser ES, Lin SP. Schizophrenia after prenatal exposure to the Dutch Hunger Winter of 1944–1945. *Arch Gen Psychiatry*. 1992;49:983–988.
- Susser E, Neugebauer R, Hoek HW, et al. Schizophrenia after prenatal famine. Further evidence. *Arch Gen Psychiatry*. 1996;53:25–31.
- St Clair D, Xu M, Wang P, et al. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959–1961. *JAMA*. 2005;294:557–562.
- Xu M, Wen-Sheng S, Liu B, et al. Prenatal malnutrition and adult schizophrenia: further evidence from the 1959–1961 famine. *Schizophr Bull*. 2009;35:568–576.
- Holick MF. Environmental factors that influence the cutaneous production of vitamin D. *Am J Clin Nutr*. 1995;61(3 suppl):638S–645S.
- Webb AR, Kline L, Holick MF. Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin. *J Clin Endocrinol Metab*. 1988;67:373–378.
- Calvo MS, Whiting SJ, Barton CN. Vitamin D intake: a global perspective of current status. *J Nutr*. 2005;135:310–316.
- Eyles D, Brown J, Mackay-Sim A, McGrath J, Feron F. Vitamin D3 and brain development. *Neuroscience*. 2003;118:641–653.
- Ko P, Burkert R, McGrath J, Eyles D. Maternal vitamin D(3) deprivation and the regulation of apoptosis and cell cycle during rat brain development. *Brain Res Dev Brain Res*. 2004;153(1):61–68.
- Cui X, McGrath JJ, Burne TH, Mackay-Sim A, Eyles DW. Maternal vitamin D depletion alters neurogenesis in the developing rat brain. *Int J Dev Neurosci*. 2007;25:227–232.
- Feron F, Burne TH, Brown J, et al. Developmental Vitamin D3 deficiency alters the adult rat brain. *Brain Res Bull*. 2005;65(2):141–148.
- Burne TH, Becker A, Brown J, Eyles DW, Mackay-Sim A, McGrath JJ. Transient prenatal vitamin D deficiency is associated with hyperlocomotion in adult rats. *Behav Brain Res*. 2004;154:549–555.
- Kesby JP, Burne TH, McGrath JJ, Eyles DW. Developmental vitamin D deficiency alters MK 801-induced hyperlocomotion in the adult rat: an animal model of schizophrenia. *Biol Psychiatry*. 2006;60:591–596.
- O'Loan J, Eyles DW, Kesby J, Ko P, McGrath JJ, Burne TH. Vitamin D deficiency during various stages of

- pregnancy in the rat; its impact on development and behaviour in adult offspring. *Psychoneuroendocrinology*. 2007;32: 227–234.
21. Kesby JP, Cui X, O'Loan J, McGrath JJ, Burne TH, Eyles DW. Developmental vitamin D deficiency alters dopamine-mediated behaviors and dopamine transporter function in adult female rats. *Psychopharmacology (Berl)*. 2010;208: 159–168.
22. Becker A, Eyles DW, McGrath JJ, Grecksch G. Transient prenatal vitamin D deficiency is associated with subtle alterations in learning and memory functions in adult rats. *Behav Brain Res*. 2005;161:306–312.
23. Abreu DA, Nivet E, Baril N, Khrestchatisky M, Roman F, Feron F. Developmental vitamin D deficiency alters learning in C57BL/6J mice. *Behav Brain Res*. 2010;208:603–608.
24. Grecksch G, Ruthrich H, Holtt V, Becker A. Transient prenatal vitamin D deficiency is associated with changes of synaptic plasticity in the dentate gyrus in adult rats. *Psychoneuroendocrinology*. 2009;34(suppl 1):S258–S264.
25. Holick MF. Deficiency of sunlight and vitamin D. *BMJ*. 2008;336:1318–1319.
26. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr*. 2008; 87:1080S–1086S.
27. Hollis BW, Wagner CL. Vitamin D deficiency during pregnancy: an ongoing epidemic. *Am J Clin Nutr*. 2006; 84:273.
28. McGrath J, Burne TH, Feron F, Mackay-Sim A, Eyles DW. Developmental vitamin D deficiency and risk of schizophrenia: a 10-year update. *Schizophr Bull*. September 10, 2010; doi:10.1093/schbul/sbq101.
29. Torrey EF, Miller J, Rawlings R, Yolken RH. Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Schizophr Res*. 1997;28:1–38.
30. Davies G, Welham J, Chant D, Torrey EF, McGrath J. A systematic review and meta-analysis of Northern Hemisphere season of birth studies in schizophrenia. *Schizophr Bull*. 2003;29:587–593.
31. Kinney DK, Teixeira P, Hsu D, et al. Relation of schizophrenia prevalence to latitude, climate, fish consumption, infant mortality, and skin color: a role for prenatal vitamin d deficiency and infections? *Schizophr Bull*. 2009; 35:582–595.
32. Saha S, Chant DC, Welham JL, McGrath JJ. The incidence and prevalence of schizophrenia varies with latitude. *Acta Psychiatr Scand*. 2006;114:36–39.
33. Dealberto MJ. Ethnic origin and increased risk for schizophrenia in immigrants to recent and traditional countries of immigration. *Acta Psychiatr Scand*. 2010;121:325–339.
34. Cantor-Graae E, Selten JP. Schizophrenia and migration: a meta-analysis and review. *Am J Psychiatry*. 2005;162: 12–24.
35. Kirkbride JB, Jones PB. *Putative Prevention Strategies to Reduce Serious Mental Illness in Migrant and Black and Minority Ethnic Groups. Foresight Mental Capital and Well-being: Discussion Paper 12*. London: Her Majesty's Stationary Office; 2008.
36. McGrath J, Eyles D, Mowry B, Yolken R, Buka S. Low maternal vitamin D as a risk factor for schizophrenia: a pilot study using banked sera. *Schizophr Res*. 2003;63:73–78.
37. McGrath J, Saari K, Hakko H, et al. Vitamin D supplementation during the first year of life and risk of schizophrenia: a Finnish birth-cohort study. *Schizophr Res*. 2004;67:237–245.
38. McGrath JJ, Eyles DW, Pedersen CB, et al. Neonatal vitamin D status and risk of schizophrenia: a population-based case-control study. *Arch Gen Psychiatry*. 2010;67:889–894.
39. Susser E, Brown A, Klonowski E, Allen R, Lindenbaum J. Schizophrenia and impaired homocysteine metabolism: a possible association. *Biol Psychiatry*. 1998;44:141–143.
40. Wolff T, Witkop CT, Miller T, Syed SB. Folic acid supplementation for the prevention of neural tube defects: an update of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2009;150:632–639.
41. Lipton SA, Kim WK, Choi YB, et al. Neurotoxicity associated with dual actions of homocysteine at the N-methyl-D-aspartate receptor. *Proc Natl Acad Sci U S A*. 1997;94: 5923–5928.
42. Hofmann MA, Kohl B, Zumbach MS, et al. Hyperhomocyst(e)inemia and endothelial dysfunction in IDDM. *Diabetes Care*. 1998;21:841–848.
43. Kamudhamas A, Pang L, Smith SD, Sadovsky Y, Nelson DM. Homocysteine thiolactone induces apoptosis in cultured human trophoblasts: a mechanism for homocysteine-mediated placental dysfunction? *Am J Obstet Gynecol*. 2004;191:563–571.
44. Picker J, Coyle J. Do maternal folate and homocysteine levels play a role in neurodevelopmental processes that increase risk for schizophrenia? *Harv Rev Psychiatry*. 2005;13:197–205.
45. Brown A, Bottiglieri T, Schaefer C, et al. Elevated prenatal homocysteine levels as a risk factor for schizophrenia. *Arch Gen Psychiatry*. 2007;64:31–39.
46. WHO. *The Prevalence of Anaemia in Women: A Tabulation of Available Information*. 2nd ed. Geneva, Switzerland: World Health Organization; 1992.
47. Crompton DW, Nesheim MC. Nutritional impact of intestinal helminthiasis during the human life cycle. *Annu Rev Nutr*. 2002;22:35–59.
48. Prentice AM. Iron metabolism, malaria, and other infections: what is all the fuss about? *J Nutr*. 2008;138:2537–2541.
49. McLean E, Egli I, Cogswell M, De Benoist B, Wojdyla D. Worldwide prevalence of anemia in preschool age children, pregnant women and non-pregnant women of reproductive age. In: Kraemer K, Zimmermann M, eds. *Nutritional Anemia*. Vol 1. Basel, Switzerland: Sight and Life Press; 2007:1–12.
50. Beard JL, Connor JR. Iron status and neural functioning. *Annu Rev Nutr*. 2003;23:41–58.
51. Laruelle M, Abi-Dargham A, van Dyck CH, et al. Single photon emission computerized tomography imaging of amphetamine-induced dopamine release in drug-free schizophrenic subjects. *Proc Natl Acad Sci U S A*. 1996; 93:9235–9240.
52. Akil M, Pierri JN, Whitehead RE, et al. Lamina-specific alterations in the dopamine innervation of the prefrontal cortex in schizophrenic subjects. *Am J Psychiatry*. 1999; 156:1580–1589.
53. Meyer-Lindenberg A, Miletich RS, Kohn PD, et al. Reduced prefrontal activity predicts exaggerated striatal dopaminergic function in schizophrenia. *Nat Neurosci*. 2002; 5:267–271.
54. Connor J, Menzies S. Relationship of iron to oligodendrocytes and myelination. *Glia*. 1996;17:83–93.
55. Kwik-Urbe CL, Gietzen D, German JB, Golub MS, Keen CL. Chronic marginal iron intakes during early development in mice result in persistent changes in dopamine metabolism and myelin composition. *J Nutr*. 2000;130:2821–2830.

56. Ortiz E, Pasquini JM, Thompson K, et al. Effect of manipulation of iron storage, transport, or availability on myelin composition and brain iron content in three different animal models. *J Neurosci Res*. 2004;77:681–689.
57. Davis KL, Stewart DG, Friedman JI, et al. White matter changes in schizophrenia: evidence for myelin-related dysfunction. *Arch Gen Psychiatry*. 2003;60:443–456.
58. Hakak Y, Walker JR, Li C, et al. Genome-wide expression analysis reveals dysregulation of myelination-related genes in chronic schizophrenia. *Proc Natl Acad Sci U S A*. 2001;98:4746–4751.
59. Flynn SW, Lang DJ, Mackay AL, et al. Abnormalities of myelination in schizophrenia detected in vivo with MRI, and post-mortem with analysis of oligodendrocyte proteins. *Mol Psychiatry*. 2003;8:811–820.
60. Carlson ES, Stead JD, Neal CR, Petryk A, Georgieff MK. Perinatal iron deficiency results in altered developmental expression of genes mediating energy metabolism and neuronal morphogenesis in hippocampus. *Hippocampus*. 2007;17:679–691.
61. Bourque SL, Iqbal U, Reynolds JN, Adams MA, Nakatsu K. Perinatal iron deficiency affects locomotor behavior and water maze performance in adult male and female rats. *J Nutr*. 2008;138:931–937.
62. Ranade SC, Rose A, Rao M, Gallego J, Gressens P, Mani S. Different types of nutritional deficiencies affect different domains of spatial memory function checked in a radial arm maze. *Neuroscience*. 2008;152:859–866.
63. McEchron MD, Goletiani CJ, Alexander DN. Perinatal nutritional iron deficiency impairs noradrenergic-mediated synaptic efficacy in the CA1 area of rat hippocampus. *J Nutr*. 2010;140:642–647.
64. Scholl TO, Hediger ML. Anemia and iron-deficiency anemia: compilation of data on pregnancy outcome. *Am J Clin Nutr*. 1994;59(2 suppl):492S–500S.
65. Kunugi H, Nanko S, Murray RM. Obstetric complications and schizophrenia: prenatal underdevelopment and subsequent neurodevelopmental impairment. *Br J Psychiatry Suppl*. 2001;40:s25–s29.
66. Cannon TD, Bearden CE, Hollister JM, Rosso IM, Sanchez LE, Hadley T. Childhood cognitive functioning in schizophrenia patients and their unaffected siblings: a prospective cohort study. *Schizophr Bull*. 2000;26:379–393.
67. Zornberg GL, Buka SL, Tsuang MT. Hypoxic-ischemia-related fetal/neonatal complications and risk of schizophrenia and other nonaffective psychoses: a 19-year longitudinal study. *Am J Psychiatry*. 2000;157:196–202.
68. Dalman C, Thomas HV, David AS, Gentz J, Lewis G, Allebeck P. Signs of asphyxia at birth and risk of schizophrenia—population-based case-control study. *Br J Psychiatry*. 2001;179:403–408.
69. Walter T, Andraca I, Chadud P, Perales C. Iron deficiency anemia: adverse effects on infant psychomotor development. *Pediatrics*. 1989;84:7–17.
70. Deinard A, Gilbert A, Dodds M, Egeland B. Iron deficiency and behavioral deficits. *Pediatrics*. 1981;68:828–833.
71. Lozoff B, Beard J, Connor J, Felt B, Georgieff M, Schallert T. Long-lasting neural and behavioral effects of iron deficiency in infancy. *Nutr Rev*. 2006;64:S34–S43.
72. Schiffman J, Sorensen H, Maeda J, et al. Childhood motor coordination and adult schizophrenia spectrum disorders. *Am J Psychiatry*. 2009;166:1041–1047.
73. Fuller R, Nopoulos P, Arndt S, O’Leary D, Ho B, Andreasen NC. Longitudinal assessment of premorbid cognitive functioning in patients with schizophrenia through examination of standardized scholastic test performance. *Am J Psychiatry*. 2002;159:1183–1189.
74. Jones P, Rodgers B, Murray R, Marmot M. Child developmental risk factors for adult schizophrenia in the British 1946 birth cohort. *Lancet*. 1994;344:1398–1402.
75. Insel B, Schaefer C, McKeague I, Susser E, Brown A. Maternal iron deficiency and the risk of schizophrenia in offspring. *Arch Gen Psychiatry*. 2008;65:1136–1144.
76. Sorensen HJ, Nielsen PR, Pedersen CB, Mortensen PB. Association between prepartum maternal iron deficiency and offspring risk of schizophrenia: population-based cohort study with linkage of Danish National Registers. January 21, 2010; doi:10.1093/schbul/sbp167.
77. Bodnar LM, Catov JM, Zmuda JM, et al. Maternal serum 25-hydroxyvitamin D concentrations are associated with small-for-gestational age births in white women. *J Nutr*. 2010;140:999–1006.
78. McGrath J. Is it time to trial vitamin D supplements for the prevention of schizophrenia? *Acta Psychiatr Scand*. 2010;121:321–324.
79. Eyles DW, Feron F, Cui X, et al. Developmental vitamin D deficiency causes abnormal brain development [first published on line September 24, 2010]. *Psychoneuroendocrinology*. doi:10.1016/j.neulet.2010.09.057.
80. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357:266–281.
81. Hypponen E, Boucher BJ. Avoidance of vitamin D deficiency in pregnancy in the United Kingdom: the case for a unified approach in National policy. *Br J Nutr*. 2010;104:1–6.
82. International Nutritional Anemia Consultative Group. *Guidelines for the eradication of iron deficiency anaemia. A report of the International Nutritional Anaemia Consultative Group*. Washington, DC: The Nutrition Foundation; 1977.
83. Mahomed K. Iron supplementation in pregnancy. *Cochrane Database Syst Rev*. 2000;(2):CD000117.
84. Villar J, Merialdi M, Gulmezoglu AM, et al. Nutritional interventions during pregnancy for the prevention or treatment of maternal morbidity and preterm delivery: an overview of randomized controlled trials. *J Nutr*. 2003;133(suppl 2):1606S–1625S.
85. Pena-Rosas JP, Viteri FE. Effects and safety of preventive oral iron or iron+folic acid supplementation for women during pregnancy. *Cochrane Database Syst Rev*. 2009;(4):CD004736.
86. Christian P, West KP, Khatry SK, et al. Effects of maternal micronutrient supplementation on fetal loss and infant mortality: a cluster-randomized trial in Nepal. *Am J Clin Nutr*. 2003;78:1194–1202.
87. Taylor DJ, Mallen C, McDougall N, Lind T. Effect of iron supplementation on serum ferritin levels during and after pregnancy. *BJOG*. 1982;89:1011–1017.
88. Szajewska H, Ruszczynski M, Chmielewska A. Effects of iron supplementation in nonanemic pregnant women, infants, and young children on the mental performance and psychomotor development of children: a systematic review of randomized controlled trials. *Am J Clin Nutr*. 2010;91:1684–1690.
89. Zhou SJ, Gibson RA, Crowther CA, Baghurst P, Makrides M. Effect of iron supplementation during pregnancy on the intelligence quotient and behavior of children at 4 y of age:

- long-term follow-up of a randomized controlled trial. *Am J Clin Nutr.* 2006;83:1112–1117.
90. Parsons AG, Zhou SJ, Spurrier NJ, Makrides M. Effect of iron supplementation during pregnancy on the behaviour of children at early school age: long-term follow-up of a randomised controlled trial. *Br J Nutr.* 2008;99:1133–1139.
91. WHO. *Standards for Maternal and Neonatal Care. Integrated Management of Pregnancy and Childbirth (IMPAC). Vol 1.8. Geneva, Switzerland: DoMPS; 2006:1–6.*
92. Khalafallah A, Dennis A, Bates J, et al. A prospective randomized, controlled trial of intravenous versus oral iron for moderate iron deficiency anaemia of pregnancy. *J Intern Med.* 2010;268:286–295.
93. Milman N. Iron prophylaxis in pregnancy—general or individual and in which dose? *Ann Hematol.* 2006;85:821–828.
94. Hotez PJ, Fenwick A, Savioli L, Molyneux DH. Rescuing the bottom billion through control of neglected tropical diseases. *Lancet.* 2009;373:1570–1575.
95. Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Ehrlich Sachs S, Sachs JD. Incorporating a rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and malaria. *PLoS Med.* 2006;3:e102.
96. Czeizel AE, Dobo M, Vargha P. Hungarian cohort-controlled trial of periconceptional multivitamin supplementation shows a reduction in certain congenital abnormalities. *Birth Defects Res A Clin Mol Teratol.* 2004;70:853–861.
97. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med.* 1992;327:1832–1835.
98. (CDC) CfDCaP. Spina bifida and anencephaly before and after folic acid mandate, United States, 1995–1996 and 1999–2000. *MMWR Morb Mortal Wkly Rep.* 2004;53:362–365.
99. De Wals P, Tairou F, Van Allen MI, et al. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med.* 2007;357:135–142.
100. Canfield MA, Collins JS, Botto LD, et al. Changes in the birth prevalence of selected birth defects after grain fortification with folic acid in the United States: findings from a multi-state population-based study. *Birth Defects Res A Clin Mol Teratol.* 2005;73:679–689.
101. Bell KN, Oakley GP, Jr. Update on prevention of folic acid-preventable spina bifida and anencephaly. *Birth Defects Res A Clin Mol Teratol.* 2009;85:102–107.
102. (FDA) UFAaD. Food standards: amendments of standards of identity for enriched grain to require addition of folic acid. *Fed Regist.* 1996;61:8781–8807.
103. Daly S, Mills JL, Molloy AM, et al. Minimum effective dose of folic acid for food fortification to prevent neural-tube defects. *Lancet.* 1997;350:1666–1669.
104. Wald NJ, Law MR, Morris JK, Wald DS. Quantifying the effect of folic acid. *Lancet.* 2001;358:2069–2073.
105. Veena SR, Krishnaveni GV, Srinivasan K, et al. Higher maternal plasma folate but not vitamin B-12 concentrations during pregnancy are associated with better cognitive function scores in 9- to 10- year-old children in South India. *J Nutr.* 2010;140:1014–1022.
106. Tamura T, Goldenberg RL, Chapman VR, Johnston KE, Ramey SL, Nelson KG. Folate status of mothers during pregnancy and mental and psychomotor development of their children at five years of age. *Pediatrics.* 2005;116:703–708.
107. Julvez J, Fortuny J, Mendez M, Torrent M, Ribas-Fito N, Sunyer J. Maternal use of folic acid supplements during pregnancy and four-year-old neurodevelopment in a population-based birth cohort. *Paediatr Perinat Epidemiol.* 2009;23:199–206.
108. del Rio Garcia C, Torres-Sanchez L, Chen J, et al. Maternal MTHFR 677C>T genotype and dietary intake of folate and vitamin B(12): their impact on child neurodevelopment. *Nutr Neurosci.* 2009;12(1):13–20.
109. Duthie SJ. Folate and cancer: how DNA damage, repair and methylation impact on colon carcinogenesis [published online ahead of print June 11, 2010]. *J Inherit Metab Dis.* doi:10.1007/s10545-010-9128-0.
110. Sanjoaquin MA, Allen N, Couto E, Roddam AW, Key TJ. Folate intake and colorectal cancer risk: a meta-analytical approach. *Int J Cancer.* 2005;113:825–828.
111. Cole BF, Baron JA, Sandler RS, et al. Folic acid for the prevention of colorectal adenomas: a randomized clinical trial. *JAMA.* 2007;297:2351–2359.
112. Mason JB, Dickstein A, Jacques PF, et al. A temporal association between folic acid fortification and an increase in colorectal cancer rates may be illuminating important biological principles: a hypothesis. *Cancer Epidemiol Biomarkers Prev.* 2007;16:1325–1329.
113. Song J, Sohn KJ, Medline A, Ash C, Gallinger S, Kim YI. Chemopreventive effects of dietary folate on intestinal polyps in Apc+/-Msh2-/- mice. *Cancer Res.* 2000;60:3191–3199.
114. Duthie SJ. Folic-acid-mediated inhibition of human colon-cancer cell growth. *Nutrition.* 2001;17:736–737.
115. Johnston RB Jr. Will increasing folic acid in fortified grain products further reduce neural tube defects without causing harm?: consideration of the evidence. *Pediatr Res.* 2008;63:2–8.
116. Steegers-Theunissen RP, Obermann-Borst SA, Kremer D, et al. Periconceptional maternal folic acid use of 400 microg per day is related to increased methylation of the IGF2 gene in the very young child. *PLoS One.* 2009;4:e7845.
117. Tam C, McKenna K, Goh YI, et al. Periconceptional folic acid supplementation: a new indication for therapeutic drug monitoring. *Ther Drug Monit.* 2009;31:319–326.
118. Troen AM, Mitchell B, Sorensen B, et al. Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr.* 2006;136:189–194.
119. Whitrow MJ, Moore VM, Rumbold AR, Davies MJ. Effect of supplemental folic acid in pregnancy on childhood asthma: a prospective birth cohort study. *Am J Epidemiol.* 2009;170:1486–1493.
120. Haberg SE, London SJ, Stigum H, Nafstad P, Nystad W. Folic acid supplements in pregnancy and early childhood respiratory health. *Arch Dis Child.* 2009;94:180–184.
121. Brown AS, Susser ES. Homocysteine and schizophrenia: from prenatal to adult life. *Progress Neuropsychopharmacol Biol Psychiatry.* 2005;29:1175–1180.
122. Ronningen KS, Paltiel L, Meltzer HM, et al. The biobank of the Norwegian Mother and Child Cohort Study: a resource for the next 100 years. *Eur J Epidemiol.* 2006;21:619–625.
123. Landrigan PJ, Trasande L, Thorpe LE, et al. The National Children's Study: a 21-year prospective study of 100,000 American children. *Pediatrics.* 2006;118:2173–2186.
124. McGrath JJ, Saha S, Burne TH, Eyles DW. A systematic review of the association between common single nucleotide

- polymorphisms and 25-hydroxyvitamin D concentrations. *J Steroid Biochem Mol Biol*. 2010;121:471–477.
125. Miyaki K. Genetic polymorphisms in homocysteine metabolism and response to folate intake: a comprehensive strategy to elucidate useful genetic information. *J Epidemiol*. 2010; 20:266–270.
126. Benyamin B, McRae AF, Zhu G, et al. Variants in TF and HFE explain approximately 40% of genetic variation in serum-transferrin levels. *Am J Hum Genet*. 2009;84:60–65.
127. van Os J, Rutten BP, Poulton R. Gene-environment interactions in schizophrenia: review of epidemiological findings and future directions. *Schizophr Bull*. 2008;34:1066–1082.